

Ten Statistical/Quantitative

Issues for Improved Environmental Regulation of Chemicals

Lauren Zeise

Office of Environmental Health Hazard
Assessment

April 29, 2004







Development of default cancer risk assessment methodologies that address potential early in life susceptibility

- Health and Safety Code, Section 901


Rodent Bioassays: Dosing Periods and Critical Windows




Tumors of nervous system in rats treated with N-ethylnitrosourea (ENU) at different ages (Naito et al., 1981)



**Total
nerve
tumors**




Effect of age at exposure on angiosarcomas induced by equal exposures (100-hr) to Vinyl Chloride



Cal/EPA's Age Related Cancer Susceptibility (“ARCS”) Database


- Cancer studies with early life vs. adult exposures
- Analyze age-at-exposure effects for a variety of chemical carcinogens
- Develop risk methods to account for fetal, infant, and pre- and post- pubescent childhood exposures
 - Default “dose effectiveness” factors or functions
 - More detailed modeling approaches for robust data sets

Default Factor



Cancer Risk = (cancer potency) x (exposure)

Function for age-dependent changes in cancer potency



Issue #1: Addressing cancer potencies with finite probability of being zero


Model

$$p = 1 - \exp[-(q_0 + q_1 d + q_2 d^2 + \dots)]$$

Potency = q_1
(statistical distribution)

Dose effectiveness:

$$q_{1 \text{ young}} / q_{1 \text{ adult}}$$



Issue #3: Adjustment for studies of same design but for age elapsed time between dosing and observation

Standard default assumption:


$$q(T) = q(T_e) \cdot (T/T_e)^3$$

(NTP “poly-3”, CalEPA cancer potency analyses, Gold’s TD50 derivation)

Observed values vary with tumor site in animals and humans.

Issue #4: Risks when agent causes cancer at multiple sites



Tumor site	Average Dose ¹ (mg/kg day)				Trend ²
	8	7	14	21	
Clitoral gland carcinomas or adenomas	3/18	27/44	48/74	41/51	$p < 0.001$
Zyodot's gland carcinomas or adenomas	1/60	12/43	21/74	16/59	$p < 0.001$
Mammary gland adenocarcinomas	1/60	2/43	14/73	20/57	$p < 0.001$
Vaginal basal cell carcinomas or adenomas	0/18	4/44	3/48	2/51	$p = 0.13$
Oval cavity squamous cell carcinomas or papillomas	2/60	2/43	6/68	5/52	$p = 0.068$
Large intestine adenocarcinomas or adenomatous polyps	0/18	1/44	1/48	3/51	$p = 0.017$
Hepatocellular carcinomas or neoplastic nodules	0/18	1/44	0/47	3/58	$p = 0.017$
Uterocervix adenoma or carcinoma	0/18	4/44	2/48	2/51	$p = 0.18$



Dimethoxybenzidine

Needed: Tools to calculate lower bounds on e.g. ED01 and ED10

Issue #5. Site specificity of mechanistic/pharmacokinetic analyses – sensitive site in humans may be overlooked.



Model: series of differential equations:

$$\frac{dAMT_{at}}{dt} = Dose \cdot Q_{in} + \frac{AMT_{lung}}{V_{lung}} \cdot \frac{Q_{out}}{P_{at}} \cdot P_{atm} - \frac{AMT_{at}}{V_{at}} \cdot Q_{out} \cdot P_{atm} - \frac{AMT_{at}}{V_{at}} \cdot Q_{in}$$

Napthalene TR-410

Issue #6. Procedures for Validating Models Applied for Regulatory Purposes

A: 2-stage model



B: 3-stage model




C: 4-stage model



D: 5-stage model





7. Creating tools to translate biological data on **variability** into mathematical models of human cancer risk

For example from NIEHS'


Environmental Genome Project

National Center for Toxicogenomics

Issue #8. Risk estimates for non-cancer endpoints

- Methyl mercury
- Arsenic
- Lead
- DBCP
- Dioxins

Issue #9. Background exposures to “complete carcinogens” and chemicals that act at selected stages




Issue #10: New systems for evaluating evidence of hazard

Too many chemicals, too little time
and money

Example



The polybrominated dipenylethers (PBDEs)




- Added to many consumer products
- Flame retardant
- Saves lives






Recognition of rising PBDE levels . . .

Organohalogen Compounds in Human Milk in Sweden (Norén and Mieronyté, 1998)





PBDEs in Herring Gull Eggs - Great Lakes




Graph from
M. Alaee
(2002)

PBDEs in U.S.


Women




Structural Similarity of PBDEs, Their Metabolites and Environmental Derivatives to T4 and PCBs




2,2',4,4',5,5'-hexachlorobiphenyl
(PCB-153)




Thyroxine (T4)




2,2',4,4',6-pentabromodiphenylether
(PBDE-100)



2-(2',4'-dibromophenoxy)-4,6-dibromoanisole
(methoxy-PBDE)



4-(2',4',6'-tribromophenoxy)-2,6-dibromophenol
(hydroxy-PBDE)



2,3,4,7,8-pentabromodibenzofuran
(PBDF)

California Ban by Legislature (Assembly Bill 302)

“ On and after January 1, 2008, a person may not manufacture, process, or distribute in commerce a product, or a flame-retarded part of a product, containing more than one-tenth of 1 percent of pentaBDE or octaBDE, by mass.”

Further tiering of regulation and evidence systems?

- Greater use of structure activity, test batteries to establish probable carcinogenicity
- Quantitative indices for “possible” that would trigger action
- Evidence for probable toxicity triggering action (addition to, e.g., hazardous air pollutants lists)
- Intermediate regulatory response for suspicious “possible” classes of chemicals